

SPECIALTY UPDATE

What's New in Limb Lengthening and Deformity Correction

Reggie C. Hamdy, MD, MSc, FRCS(C), Mitchell A. Bernstein, MD, FRCS(C), Austin T. Fragomen, MD, and S. Robert Rozbruch, MD

Limb reconstruction surgery is an emerging subspecialty. In the past year, 5 major events highlighted the growing interest in this field. The first combined meeting of the 3 international societies devoted to the study and advancement of limb lengthening and reconstruction—the Association for the Study and Application of the Methods of Ilizarov (ASAMI) International, the Limb Lengthening and Reconstruction Society (LLRS)-North America, and the International LLRS (ILLRS)—was held in November 2015, in Miami. Second, a new journal, the *Journal of Limb Lengthening & Reconstruction*, was launched. Third, the *Limb Lengthening and Reconstruction Surgery Case Atlas* was published (Springer International; S.R.R. and R.C.H., editors). This is a major work in association with the LLRS, with 3 volumes and 302 cases covering all aspects of limb reconstruction. Fourth, the LLRS supported and participated in the AAOS/OREF/ORS (American Academy of Orthopaedic Surgeons/Orthopaedic Research and Education Foundation/Orthopaedic Research Society) Clinician Scholar Career Development Program. Lastly, the LLRS Traveling Fellowship program was implemented.

In addition to these events, numerous articles related to limb reconstruction and deformity correction were published in 2015. This Specialty Update provides a summary of this recent research.

Pediatrics

Congenital Limb Deficiencies

Several studies investigated the treatment of congenital limb deficiencies. Shahcheraghi and Javid found that the majority of children with tibial hemimelia had a good functional outcome following reconstructive procedures or knee disarticulation¹. In

a study involving the largest series of tibial hemimelia cases reported from a single institution, other investigators used a modified Jones classification that included a “type V” for cases with global deficiency of the tibia. There were no cases of type III². In a study involving children >4 years of age with type-II fibular hemimelia, Popkov et al. concluded that tibial lengthening should be associated with resection of the fibular anlage in order to prevent recurrence of the valgus deformity of the ankle³. On the basis of radiographic assessment of the feet of patients with fibular hemimelia, Reyes et al. noted that the deficiency “should not be viewed as a global ‘lateral lower-limb deficiency’ nor the foot ray deficiency as ‘lateral.’”⁴ Other investigators found satisfactory results following a specific protocol of lengthening in children with congenital femoral deficiency⁵. Osteonecrosis of the femoral head in the contralateral hip was described for 3 children with congenital femoral deficiency⁶. Prophylactic femoral rodding with Rush rods following fixator removal was reported to be a safe and effective method to decrease the incidence of fracture following lengthening⁷. However, femoral lengthening with circular external fixation had a higher rate of complications than that with use of a motorized nail⁸.

Congenital Pseudarthrosis of the Tibia (CPT)

Three studies provided additional insight into the treatment of CPT in pediatric patients. Comparable results were found between those with and without neurofibromatosis type I who were treated with use of the same Ilizarov technique⁹. Refracture was the main issue following Ilizarov bone transport¹⁰. Proximal tibial lengthening after initial union of CPT was an effective form of management of tibial shortening¹¹.

Blount Disease

For patients with Blount disease who require minimal lengthening, correction using the Taylor Spatial Frame (Smith & Nephew)

Specialty Update has been developed in collaboration with the Board of Specialty Societies (BOS) of the American Academy of Orthopaedic Surgeons.

Disclosure: One or more of the authors received a stipend from JBJS for writing this work. On the **Disclosure of Potential Conflicts of Interest** forms, which are provided with the online version of the article, one or more of the authors checked “yes” to indicate that the author had a relevant financial relationship in the biomedical arena outside the submitted work.

WHAT'S NEW IN LIMB LENGTHENING AND DEFORMITY CORRECTION

can potentially be performed safely without osteotomy and fixation of the fibula¹². A new classification system for complications in patients with Blount disease treated with a circular external fixator also was presented¹³.

Skeletal Dysplasia

Several reports addressed lengthening in children with achondroplasia. First, tibial lengthening had a significantly lower complication rate than did femoral lengthening¹⁴. Second, although a complication rate of 70% was reported for children with achondroplasia who underwent lengthening procedures, none of these children were left with any long-term sequelae, and all were pleased with the results¹⁵. Third, although double-level tibial lengthening can be difficult, the results are usually gratifying¹⁶. Finally, humeral lengthening was reported to increase the independence of children with achondroplasia, indicating that it is more than a cosmetic procedure¹⁷.

In addition, the successful use of the Fassier-Duval telescopic rod in the correction of deformity in 4 patients with Stüve-Wiedemann syndrome was reported¹⁸. A treatment protocol for the correction of deformity in patients with skeletal dysplasia involving the use of a circular external fixator was also presented¹⁹.

New Techniques

In children with osteogenesis imperfecta, the nonunion of long bones was successfully treated with compressed sandwich allograft cortical struts²⁰, and unicortical locking-plate fixation was successfully applied as an adjunct to intramedullary rodding in cases of fractures and osteotomies²¹. Ippolito et al. reported on a 2-stage technique for the successful treatment of patients with polyostotic fibrous dysplasia and severe coxa vara²². The use of flexible intramedullary nails augmented by external fixation as an alternative treatment of open metaphyseal diaphyseal tibial fractures in adolescents was reported by Atef and Tantawy, with satisfactory results²³.

A new technique for the resection of a central or mixed physeal bar by a complete transverse osteotomy at the metaphysis was also reported²⁴. In another report, satisfactory results of plate-assisted lengthening of the femur and tibia were found²⁵. No improvement in function was found for 11 patients with Legg-Calvé-Perthes disease or developmental dysplasia of the hip who underwent femoral head reduction osteotomy²⁶. Other investigators described a novel technique for the treatment of chronic patellar sleeve fractures using the Taylor Spatial Frame²⁷. Laine et al. reported on the management of 8 patients with open fractures of the tibia with bone loss²⁸. The technique included static stabilization for soft-tissue healing and acute shortening, with a plan for subsequent limb lengthening. Ulnar osteotomy and gradual correction using the Ilizarov external fixator was described for 4 cases of chronic anterior Monteggia lesion²⁹. A successful modified technique for the use of the Taylor Spatial Frame in patients with cubitus

varus after malunion of a pediatric supracondylar fracture was reported³⁰. The risk of proximal rod migration in patients with osteogenesis imperfecta who had deformity correction with use of a Bailey-Dubow rod could be reduced by complete correction of the angular deformity and optimal placement of the rod at the distal physis³¹. Anam et al. reviewed a series of 110 patients with osteogenesis imperfecta who had 261 intramedullary rodding procedures, 139 in the femur and 122 in the tibia, and showed that the introduction of a new medical and surgical protocol for deformity correction (a bisphosphonate infusion-free interval of 4 months after the osteotomy, use of an osteotome instead of a power saw, and the introduction of parenteral zoledronic acid) led to a decrease in the rate of delayed osteotomy healing when compared with the treatment of patients using the old regimen³². Shore et al. compared the results, cost, and complications of 42 children with diaphyseal tibial fracture treated with use of a Taylor Spatial Frame or a uniplanar fixator and reported equivalent cost for care delivery by using either fixator³³. In patients who underwent a transtibial amputation, osteocartilaginous capping with a proximal part of the ipsilateral fibula prevented recurrent overgrowth in 90% of the cases³⁴.

Other Pediatric

The combined use of internal and external osteosynthesis allowed considerable reduction in the duration of the external fixator and decreased the number of complications in children undergoing leg lengthening³⁵ and in children with X-linked hereditary hypophosphatemic rickets who underwent deformity correction³⁶. Preoperative psychological intervention as part of an interdisciplinary team approach was recommended for patients undergoing limb reconstruction³⁷. Another study sought to determine the normal radiographic values of anatomical and mechanical alignment angles of the lower limbs of children of various age groups, the first such chronological evaluation in a relatively large series of patients³⁸. As found in a cadaveric study of femora of donors 8 to 20 years of age by Park et al., nail entry at the pediatric greater trochanter apex would likely result in anterior placement³⁹. Lastly, in cases of poor or delayed bone formation during lengthening procedures, the addition of compressive forces to the distraction protocol (the accordion maneuver) may be beneficial⁴⁰.

Trauma

Hypertrophic Tibial Nonunion

Salih et al., in an attempt to predict which diaphyseal tibial fractures would progress to hypertrophic nonunion, reported on a radiographic sign, termed the *callus fracture sign*⁴¹. This had a sensitivity of 69.6%, specificity of 91.3%, and positive and negative predictive values of 88.9% and 75.0%, respectively. Xu et al. retrospectively reviewed the data of 12 patients with a diagnosis of hypertrophic nonunion treated with an Ilizarov ring fixator, with a union rate of 100%⁴². In no cases did the authors perform compression at the site of nonunion to

WHAT'S NEW IN LIMB LENGTHENING AND DEFORMITY CORRECTION

augment healing. In another study, distal tibial hypertrophic nonunions treated with a circular fixator demonstrated union in 100% of the patients⁴³. Patients treated with the Taylor Spatial Frame had improved deformity correction. In a study by Ferreira et al., the authors reported a union rate of 89.1% among 46 patients with aseptic, stiff tibial nonunions treated with hexapod external fixators and distraction⁴⁴.

Tibial Fracture

A meta-analysis compared various modalities for the treatment of open tibial shaft fractures⁴⁵. Method ranking revealed that unreamed nailing was the modality with the highest probability of being the best treatment, followed by reamed nailing, external fixation, and plate fixation. A systematic review of 57 studies evaluating the treatment of open type-III tibial shaft fractures concluded that circular frames have surgical outcomes that are equivalent to those of other techniques⁴⁶. In another systematic review evaluating the outcome of Ilizarov treatment of infected tibial and femoral nonunions, an average 97.26% rate of union was shown for the included studies⁴⁷. In a series of 56 patients treated with use of a Taylor Spatial Frame for tibial shaft fractures, the indications for circular fixation remained inconclusive⁴⁸. Circular fixation remains a powerful tool, especially for cases in which a type-IIIB injury can be converted to a type-IIIA with a technique called *intentional deformation*⁴⁹.

Lengthening and then nailing (LATN) and lengthening and then plating (LAP) are 2 methods of decreasing the time that patients spend in an external fixator. Bernstein et al. reported on 58 patients with posttraumatic bone loss (mean, 5.3 cm) treated with either classic circular fixation and bone transport or “integrated fixation” (LATN or LAP)⁵⁰. They demonstrated a 100% union rate and no difference in the severity or number of complications between the 2 groups. In a study involving an animal model of tibial fracture, there was a subclinical degeneration of myelin fibers in the peroneal and tibial nerves during fracture healing and following fixator removal⁵¹.

Distal Tibial/Pilon Fracture

In a randomized comparative study, 40 patients were treated with open reduction and internal fixation with conventional plating or Ilizarov circular fixation⁵². The rate of healing in the Ilizarov fixation group was higher ($p = 0.003$). The addition of ankle arthroscopy for visualization of the articular surface did not show additional benefit for 23 patients with closed pilon fractures treated with Ilizarov external fixation⁵³. In a retrospective case series of 28 open pilon fractures, Danoff et al. reported on their protocol of the use of staged open reduction of the articular surface and Ilizarov fixation⁵⁴.

Femoral and Tibial Bone Transport

Massive bone loss in the tibia and femur was reported in a study of 110 patients with infected nonunion treated by bone

transport⁵⁵. Ilizarov circular fixation was used in the tibia, and a monolateral rail was used in the femur. The mean bone defect, post-debridement, was 6.15 cm. The authors reported a 100% union rate, with no recurrence of infection. The successful use of a monolateral external fixator was demonstrated in another study of 7 patients with a mean infected femoral bone defect of 8.1 cm⁵⁶.

Other Trauma

The use of circular fixation with classic stainless steel Ilizarov rings has been demonstrated to be safe and cost-effective in austere and low socioeconomic settings⁵⁷⁻⁵⁹. Tibial plateau fractures, Schatzker type-IV to VI fractures, can be effectively treated with Ilizarov circular fixation⁶⁰. In a retrospective study of 177 patients, the rate of venous thromboembolism in patients with tibial fractures treated with circular fixation was 4% (deep-vein thrombosis, 1.7%, and pulmonary embolism, 2.3%)⁶¹. Zhong et al. reported on their single-stage reconstruction for severe foot and ankle deformities that included the use of a sural neurocutaneous flap, noting no flap failures or neurovascular complications⁶².

Adult

Rozbruch et al. retrospectively examined data regarding surgical cases and outpatient visits from the limb-lengthening and complex reconstruction service at one institution⁶³. Referrals were primarily from orthopaedic surgeons (56%) and self/Internet research (25%).

Limb Realignment and Arthritis

There were several studies that supported the use of realignment osteotomy to offload the arthritic compartment of a knee with osteoarthritis⁶⁴⁻⁶⁹. Stupina et al. concluded that restoring the microarchitecture of the subchondral bone and improving its vascularization after tunneling with the introduction of bone marrow suspension into the drill-holes enhances chondrocyte metabolism and recovers their functionality⁷⁰. The success of open-wedge osteotomy is impacted by the completion and speed of bone growth in the triangular space created at surgery. As found in one systematic review, the lowest rates of delayed union/nonunion were in autograft bone-filled osteotomies⁷¹. While high tibial osteotomy (HTO) is the most common knee realignment procedure performed, distal femoral osteotomy was evaluated in the treatment of 21 knees for valgus knee malalignment with arthritis or to protect a knee compartment in which chondral resurfacing surgery had been performed⁷². Harris et al. studied the outcomes of patients undergoing simultaneous meniscal transplantation, osteotomy, and articular cartilage repair, noting improvements in validated patient-reported clinical outcome scores at long-term follow-up⁷³. Konopka et al. studied the cost-effectiveness of HTO and unicompartmental knee replacement as alternatives to total knee replacement for patients 50 to 60 years of age⁷⁴.

WHAT'S NEW IN LIMB LENGTHENING AND DEFORMITY CORRECTION

Limb Lengthening

While bone lengthening with external fixation has been demonstrated to be successful, newer methods of integrated fixation are showing promise for some patients by decreasing external fixator-related complications and the duration of time needed in the frame^{75,76}. Dehghan et al. showed that the transplantation of autologous bone marrow mesenchymal stem cells (BM-MSCs) positively affected early osseous consolidation in distraction osteogenesis⁷⁷. Published results of the use of the PRECICE Intramedullary Limb Lengthening System (Ellipse Technologies) in >250 cases were reported to be excellent, with less pain and lower complication rates than with external fixation methods or previous implantable nail systems⁷⁸. Patients who had experienced both internal and external fixation preferred the internal lengthening device with respect to overall satisfaction⁷⁹. The lengthening nail might be superior to external fixation for femoral lengthening, when the anatomical conditions and the complexity of the deformity allow the use of an intramedullary nail^{80,81}. A low-cost custom knee device with polyester synthetic conformable casting material for the treatment of knee flexion contractures was used successfully in 23 patients (27 limbs) who developed knee flexion contracture during femoral lengthening with an intramedullary lengthening femoral nail⁸². Elbatrawy and Ragab studied the outcomes of 50 patients who underwent stature lengthening for psychological reasons and concluded that the Ilizarov device is a safe tool for limb lengthening in individuals of short stature⁸³.

Limb Alignment and Osseous Anatomy

Zampogna et al. compared the measurement of the anatomical femorotibial angle on a knee radiograph with the measurement of the mechanical axis on a full-length standing hip-to-ankle radiograph⁸⁴. The knee radiograph measurement had weak correlation with the long radiograph. In addition, as concluded by Lee et al., alignment after total knee arthroplasty should be assessed by measuring mechanical alignment on a full-length standing radiograph rather than by anatomical alignment⁸⁵. In another study, by Cho et al., the foot progression angle and the mechanical axis of the lower limb were significantly correlated with the first and second peak knee adduction moments⁸⁶. Weinberg et al. studied the association between rotational deformities of the femur and tibia and the development of arthritis using an osteological collection⁸⁷. As assessed by multiple regression analysis, neither tibial torsion nor femoral version proved to be independent predictors of hip or knee arthritis. Weinberg and Liu found that increasing tibial length relative to femoral length was a significant predictor of ipsilateral hip and knee arthritis⁸⁸.

Deformity Correction

A deficiency of the lateral compartment of the knee, often in the setting of skeletal dysplasia, is an intra-articular deformity resulting in genu valgum. Advancement of the lateral femoral condyle can be performed to correct the deformity⁸⁹. Alexis et al. evaluated the utility of the Taylor Spatial Frame among cases treated in Haiti⁹⁰. Good results were observed in 99% of the cases.

Symmetrical distribution of the load of the lower limbs and balance are among the determinants of proper biomechanics of the musculoskeletal system. In a study of 57 patients, load distribution of the lower limbs in the study group after realignment surgery did not significantly differ from that in normal controls⁹¹. The same group examined 56 patients who underwent derotational corticotomies of the distal epiphysis of the femur or the proximal epiphysis of the tibia using the Ilizarov method⁹². The authors found no differences between the study group and controls in physical and sports activity after surgical correction.

Limb Salvage and Amputation

Circular external fixation may be used to stabilize limbs in an intentionally short position and then gradually correct the deformity^{93,94}. The overall incidence, prevalence, fall characteristics, and risk factors were calculated for falls in an amputee population resulting in rehospitalization⁹⁵. Life-threatening infections can occur while using circular external fixation⁹⁶. A satisfactory outcome was obtained with early diagnosis and aggressive medical and surgical treatment.

Other Adult-Related

The Return to Run (RTR) clinical pathway, which was introduced in 2009, has demonstrated favorable results among returning active-duty service members in terms of running, sports participation, active duty, and deployments⁹⁷. Rivera et al. hypothesized that nonsteroidal anti-inflammatory drugs (NSAIDs) may be delivered locally in a wound for the potential prevention of heterotopic ossification⁹⁸.

Foot and Ankle

The use of fluoroscopy was recommended while using an Ilizarov semicircular external fixator for distraction osteogenesis in brachymetatarsia⁹⁹. A study of complex ankle arthrodesis showed a high fusion rate with minimal complications, confirming the utility of ringed fixators¹⁰⁰. A review of the literature on complex ankle arthrodesis yielded an algorithm for surgical treatment¹⁰¹. For a patient with complete fibular agenesis, Cavadas and Thione successfully used the proximal part of the contralateral fibula including the growth plate to craft a lateral malleolus with microvascular anastomosis¹⁰². In a study by Lohia et al. on resistant idiopathic clubfoot in toddlers, both gradual correction with external fixation and open release with acute correction were found to provide significant improvement¹⁰³. A group of plastic surgeons treated foot ulcers associated with Charcot arthropathy by several methods, including free tissue transfer, concluding that aggressive treatment can yield a high healing rate¹⁰⁴.

Additional Studies of Interest

In 4 cases of elbow arthrodesis, circular external fixation was successful at providing a stable and functional upper extremity when amputation was the alternative¹⁰⁵. Three studies showed creative uses for Ilizarov mini-fixators in the correction of hand contractures: a severe intrinsic-plus hand deformity was treated

WHAT'S NEW IN LIMB LENGTHENING AND DEFORMITY CORRECTION

with selective releases combined with fixator-assisted stretching¹⁰⁶; carpometacarpal (CMC) arthritis with associated metacarpophalangeal (MCP) joint hyperextension deformity was treated with CMC arthroplasty and metacarpal osteotomy and external fixation with excellent results¹⁰⁷; and, in the final study, postoperative rehabilitation with the use of rubber bands was emphasized¹⁰⁸. A study of 7 patients treated with resection and bone transport using a classic Ilizarov circular fixator yielded excellent or good results in all cases¹⁰⁹. A larger series of 38 transports included classic ring transport with multiple osteotomy sites and Ilizarov crossed olive-wire transport and showed excellent results for the entire group after an average lengthening of 10 cm¹¹⁰. Alzahrani et al. investigated the role of sclerostin antibody infusions in a mouse model of tibial fracture-healing and reported rapid healing when compared with controls¹¹¹. A recent review of distraction osteogenesis and methods for improving bone regeneration focused on the use of adipose stem cells. The study outlined the relative abundance of cells in the adipose tissue and the equivalent performance of these cells when compared with crest-derived cells¹¹². Improved delivery of bone morphogenetic protein (BMP) and other potent bone-healing stimulators may be possible with the use of a chitosan sponge, where the BMP is released more evenly and over a sustained period¹¹³. A case of a 10-year-old patient with anterior cruciate ligament (ACL) reconstruction-related growth arrest and subsequent deformity showed successful use of the lengthening and plating technique¹¹⁴.

In a study involving patients who had tibial surgery and external fixation, Lundblad et al. used PET/CT (positron emission tomography/computed tomography) bone scans to measure bone activity and showed that several static measurements could be used as a proxy for a complete dynamic scan, thus saving examination time¹¹⁵. An investigation of distraction osteogenesis in a sheep model focused on the callus production rate, volume, and mechanical parameters¹¹⁶. In a study of distraction osteogenesis involving dogs, investigators found that the concentrations of growth factors (GFs), including insulin-like GFs 1 and 2, stem cell factor, vascular endothelial GF, and transforming GF- β 1, varied with the rate of distraction¹¹⁷.

A novel tensioning and clamping system that resists loss of wire tension far better than traditional clamping constructs was evaluated by Bairaktari et al.¹¹⁸. Striving for dynamization, Capanni et al. found that dynamic locking plates (utilizing a silicone interface between plate and screws, providing elastically suspended locking holes) behave more like an Ilizarov fixator than a traditional locking plate¹¹⁹. Additional insight into the dynamic properties of circular fixation through sequential testing of various

frame constructs was provided by Henderson et al. Their hypothesis, that half-pins would create a cantilever system that increased the presence of undesirable strain at the fracture site, was disproved by data showing that the all-half-pin construct demonstrated less strain than the all-wire gold standard¹²⁰. Oblique fractures have presented challenges to surgeons using external fixation. Looking at this problem in detail, Jabbar and Khaleel found that placing crossed olive wires across oblique fractures resulted in the most stable constructs¹²¹. Using a transverse fracture and osteotomy model, researchers found that a hexapod-type (Ortho SUV) frame, which allowed for a dynamized mode, was better able to resist deforming forces than was a classic Ilizarov external fixator¹²².

Several other articles were dedicated to answering practical questions regarding relevant human anatomy. The method of vascularized free-tissue transfer was further refined by the use of perforating vessels in place of the larger, named vessels at the anastomosis site¹²³. The authors found no significant increase in complications, despite connecting to a reduced-flow vessel. The use of external fixation around the distal part of the femur creates concern for knee-joint contamination and septic arthritis. A cadaveric study demonstrated that the safe corridor for extra-articular fixation is narrower than perhaps previously thought¹²⁴. Iatrogenic injury of the common peroneal nerve during wire placement in the fibular head is a devastating complication of the application of circular fixation. An anatomical study was conducted among cadavers to precisely outline the safe zone for proximal fibular transfixation¹²⁵. Ktistakis et al. analyzed 369 articles on the topic of pin-site infection after external fixation and concluded that there is no clearly superior method for conducting pin care¹²⁶.

Reggie C. Hamdy, MD, MSc, FRCS(C)¹
Mitchell A. Bernstein, MD, FRCS(C)²
Austin T. Fragomen, MD³
S. Robert Rozbruch, MD³

¹Shriners Hospitals for Children-Canada, Montréal, Québec, Canada

²Loyola University Stritch School of Medicine, Maywood, Illinois

³Hospital for Special Surgery, New York, NY

E-mail address for R.C. Hamdy: rhamdy@shriners.mcgill.ca
E-mail address for M.A. Bernstein: mitchell.bernstein@lumc.edu
E-mail address A.T. Fragomen: FragomenA@HSS.edu
E-mail address S.R. Rozbruch: RozbruchSR@HSS.edu

References

1. Shahcheraghi GH, Javid M. Functional assessment in tibial hemimelia (can we also save the foot in reconstruction?). *J Pediatr Orthop*. 2015 May 14. [Epub ahead of print].
2. Clinton R, Birch JG. Congenital tibial deficiency: a 37-year experience at 1 institution. *J Pediatr Orthop*. 2015 Jun;35(4):385-90.

3. Popkov A, Aranovich A, Popkov D. Prevention of recurrence of tibia and ankle deformities after bone lengthening in children with type II fibular hemimelia. *Int Orthop*. 2015 Jul;39(7):1365-70. Epub 2015 Apr 2.
4. Reyes BA, Birch JG, Hootnick DR, Cherkashin AM, Samchukov ML. The nature of foot ray deficiency in congenital fibular deficiency. *J Pediatr Orthop*. 2015 Sep 9. [Epub ahead of print].

WHAT'S NEW IN LIMB LENGTHENING AND DEFORMITY CORRECTION

5. Prince DE, Herzenberg JE, Standard SC, Paley D. Lengthening with external fixation is effective in congenital femoral deficiency. *Clin Orthop Relat Res.* 2015 Oct;473(10):3261-71. Epub 2015 Jul 21.
6. Brown TS, Wimberly RL, Birch JG. Avascular necrosis in the contralateral hip in patients with congenital femoral deficiency: a report of 3 cases. *J Pediatr Orthop.* 2015 Nov 16. [Epub ahead of print].
7. Abdelgawad AA, Jauregui JJ, Standard SC, Paley D, Herzenberg JE. Prophylactic intramedullary rodding following femoral lengthening in congenital deficiency of the femur. *J Pediatr Orthop.* 2015 Dec 8. [Epub ahead of print].
8. Black SR, Kwon MS, Cherkashin AM, Samchukov ML, Birch JG, Jo CH. Lengthening in congenital femoral deficiency: a comparison of circular external fixation and a motorized intramedullary nail. *J Bone Joint Surg Am.* 2015 Sep 2;97(17):1432-40.
9. Borzunov DY, Chevardin AY, Mitrofanov AI. Management of congenital pseudarthrosis of the tibia with the Ilizarov method in a paediatric population: influence of aetiological factors. *Int Orthop.* 2016 Feb;40(2):331-9. Epub 2015 Nov 7.
10. Vanderstappen J, Lammens J, Berger P, Laumen A. Ilizarov bone transport as a treatment of congenital pseudarthrosis of the tibia: a long-term follow-up study. *J Child Orthop.* 2015 Aug;9(4):319-24. Epub 2015 Aug 13.
11. Zhu GH, Mei HB, He RG, Liu K, Tang J, Wu JY. Effect of distraction osteogenesis in patient with tibial shortening after initial union of congenital pseudarthrosis of the tibia (CPT): a preliminary study. *BMC Musculoskelet Disord.* 2015;16:216. Epub 2015 Aug 21.
12. Sachs O, Katzman A, Abu-Johar E, Eidelman M. Treatment of adolescent Blount disease using Taylor Spatial Frame with and without fibular osteotomy: is there any difference? *J Pediatr Orthop.* 2015 Jul-Aug;35(5):501-6.
13. Cherkashin AM, Samchukov ML, Birch JG, Da Cunha AL. Evaluation of complications of treatment of severe Blount's disease by circular external fixation using a novel classification scheme. *J Pediatr Orthop B.* 2015 Mar;24(2):123-30.
14. Park KW, Garcia RA, Rejuo CA, Choi JW, Song HR. Limb lengthening in patients with achondroplasia. *Yonsei Med J.* 2015 Nov;56(6):1656-62.
15. Donaldson J, Aftab S, Bradish C. Achondroplasia and limb lengthening: results in a UK cohort and review of the literature. *J Orthop.* 2015 Mar;12(1):31-4. Epub 2015 Jan 28.
16. Burghardt RD, Yoshino K, Kashiwagi N, Yoshino S, Bhav A, Paley D, Herzenberg JE. Bilateral double level tibial lengthening in dwarfism. *J Orthop.* 2015 Dec;12(4):242-7. Epub 2015 Jun 3.
17. Balci HI, Kocaoglu M, Sen C, Eralp L, Batbay SG, Bilsel K. Bilateral humeral lengthening in achondroplasia with unilateral external fixators: is it safe and does it improve daily life? *Bone Joint J.* 2015 Nov;97-B(11):1577-81.
18. Wright J, Kazzaz S, Hill RA. Developments in the orthopaedic management of children with Stüve-Wiedemann syndrome: use of the Fassier-Duval telescopic rod to maintain correction of deformity. *J Pediatr Orthop.* 2015 Dec 8. [Epub ahead of print].
19. Samchukov ML, Birch JG, Cherkashin AM, Makarov MR. Deformity correction in pediatric skeletal dysplasia: treatment challenges and solutions. *J Pediatr Orthop B.* 2015 Mar;24(2):131-8.
20. Puvanesarajah V, Shapiro JR, Sponseller PD. Sandwich allografts for long-bone nonunions in patients with osteogenesis imperfecta: a retrospective study. *J Bone Joint Surg Am.* 2015 Feb 18;97(4):318-25.
21. Cho TJ, Lee K, Oh CW, Park MS, Yoo WJ, Choi IH. Locking plate placement with unicortical screw fixation adjunctive to intramedullary rodding in long bones of patients with osteogenesis imperfecta. *J Bone Joint Surg Am.* 2015 May 6;97(9):733-7.
22. Ippolito E, Farsetti P, Valentini MB, Potenza V. Two-stage surgical treatment of complex femoral deformities with severe coxa vara in polyostotic fibrous dysplasia. *J Bone Joint Surg Am.* 2015 Jan 21;97(2):119-25.
23. Atef A, El Tantawy A. Open unstable metaphyseal-diaphyseal fractures of the tibia in adolescents: treatment by flexible intramedullary nails augmented by external fixator. *Int Orthop.* 2015 May;39(5):921-6. Epub 2015 Feb 19.
24. Kim HT, Lim KP, Jang JH, Ahn TY. Resection of a physeal bar with complete transverse osteotomy at the metaphysis and Ilizarov external fixation. *Bone Joint J.* 2015 Dec;97-B(12):1726-31.
25. Georgiadis AG, Rossow JK, Laine JC, Iobst CA, Dahl MT. Plate-assisted lengthening of the femur and tibia in pediatric patients. *J Pediatr Orthop.* 2015 Sep 11. [Epub ahead of print].
26. Siebenrock KA, Anwander H, Zurmühle CA, Tannast M, Slongo T, Steppacher SD. Head reduction osteotomy with additional containment surgery improves sphericity and containment and reduces pain in Legg-Calvé-Perthes disease. *Clin Orthop Relat Res.* 2015 Apr;473(4):1274-83.
27. Potini VC, Reilly MC, Gehrmann RM. Staged treatment of a chronic patellar sleeve fracture using the Taylor Spatial Frame. *Knee.* 2015 Dec;22(6):672-6. Epub 2015 May 8.
28. Laine JC, Cherkashin A, Samchukov M, Birch JG, Rathjen KE. The management of soft tissue and bone loss in type IIIB and IIIC pediatric open tibia fractures. *J Pediatr Orthop.* 2015 Apr 13. [Epub ahead of print].
29. Bor N, Rubin G, Rozen N, Herzenberg JE. Chronic anterior Monteggia lesions in children: report of 4 cases treated with closed reduction by ulnar osteotomy and external fixation. *J Pediatr Orthop.* 2015 Jan;35(1):7-10.
30. Belthur MV, Iobst CA, Bor N, Segev E, Eidelman M, Standard SC, Herzenberg JE. Correction of cubitus varus after pediatric supracondylar elbow fracture: alternative method using the Taylor Spatial Frame. *J Pediatr Orthop.* 2015 Apr 21. [Epub ahead of print].
31. Lee K, Park MS, Yoo WJ, Chung CY, Choi IH, Cho TJ. Proximal migration of femoral telescopic rod in children with osteogenesis imperfecta. *J Pediatr Orthop.* 2015 Mar;35(2):178-84.
32. Anam EA, Rauch F, Glorieux FH, Fassier F, Hamdy R. Osteotomy healing in children with osteogenesis imperfecta receiving bisphosphonate treatment. *J Bone Miner Res.* 2015 Aug;30(8):1362-8. Epub 2015 May 10.
33. Shore BJ, DiMauro JP, Spence DD, Miller PE, Glotzbecker MP, Spencer S, Hedequist D. Uniplanar versus Taylor Spatial Frame external fixation for pediatric diaphyseal tibia fractures: a comparison of cost and complications. *J Pediatr Orthop.* 2015 Jun 17. [Epub ahead of print].
34. Fedorak GT, Watts HG, Cuomo AV, Ballesteros JP, Grant HJ, Bowen RE, Scaduto AA. Osteocartilaginous transfer of the proximal part of the fibula for osseous overgrowth in children with congenital or acquired tibial amputation: surgical technique and results. *J Bone Joint Surg Am.* 2015 Apr 1;97(7):574-81.
35. Bukva B, Virgoč G, Rakovac I, Dučić S, Sindik J, Čoklo M, Marinović M, Bakota B. Complications in leg lengthening using an Ilizarov external fixator and intramedullary alignment in children: comparative study during a fourteen-year period. *Injury.* 2015 Nov;46(Suppl 6):S48-51. Epub 2015 Nov 21.
36. Popkov A, Aranovich A, Popkov D. Results of deformity correction in children with X-linked hereditary hypophosphatemic rickets by external fixation or combined technique. *Int Orthop.* 2015 Dec;39(12):2423-31. Epub 2015 Jul 7.
37. Richard HM, Nguyen DC, Birch JG, Roland SD, Samchukov MK, Cherkashin AM. Clinical implications of psychosocial factors on pediatric external fixation treatment and recommendations. *Clin Orthop Relat Res.* 2015 Oct;473(10):3154-62.
38. Popkov D, Lascombes P, Berte N, Hetzel L, Baptista BR, Popkov A, Journeau P. The normal radiological anteroposterior alignment of the lower limb in children. *Skeletal Radiol.* 2015 Feb;44(2):197-206. Epub 2014 Jul 5.
39. Park PJ, Weinberg DS, Petro KF, Liu RW. An anatomic study of the greater trochanter starting point for intramedullary nailing in the skeletally immature. *J Pediatr Orthop.* 2015 Jul 2. [Epub 2015 ahead of print].
40. Makhdom AM, Cartaleanu AS, Rendon JS, Villemure I, Hamdy RC. The accordion maneuver: a noninvasive strategy for absent or delayed callus formation in cases of limb lengthening. *Adv Orthop.* 2015;2015:912790. Epub 2015 Oct 19.
41. Salih S, Blakey C, Chan D, McGregor-Riley JC, Royston SL, Gowllett S, Moore D, Dennison MG. The callus fracture sign: a radiological predictor of progression to hypertrophic non-union in diaphyseal tibial fractures. *Strategies Trauma Limb Reconstr.* 2015 Nov;10(3):149-53. Epub 2015 Nov 24.
42. Xu J, Jia YC, Kang QL, Chai YM. Management of hypertrophic nonunion with failure of internal fixation by distraction osteogenesis. *Injury.* 2015 Oct;46(10):2030-5. Epub 2015 Jun 17.
43. Schoenleber SJ, Hutson JJ Jr. Treatment of hypertrophic distal tibia nonunion and early malunion with callus distraction. *Foot Ankle Int.* 2015 Apr;36(4):400-7. Epub 2014 Oct 30.
44. Ferreira N, Marais LC, Aldous C. Hexapod external fixator closed distraction in the management of stiff hypertrophic tibial nonunions. *Bone Joint J.* 2015 Oct;97-B(10):1417-22.
45. Foote CJ, Guyatt GH, Vignesh KN, Mundi R, Chaudhry H, Heels-Ansdell D, Thabane L, Torretta P 3rd, Bhandari M. Which surgical treatment for open tibial shaft fractures results in the fewest reoperations? A network meta-analysis. *Clin Orthop Relat Res.* 2015 Jul;473(7):2179-92. Epub 2015 Feb 28.
46. Dickson DR, Moulder E, Hadland Y, Giannoudis PV, Sharma HK. Grade 3 open tibial shaft fractures treated with a circular frame, functional outcome and systematic review of literature. *Injury.* 2015 Apr;46(4):751-8. Epub 2015 Jan 23.
47. Yin P, Ji Q, Li T, Li J, Li Z, Liu J, Wang G, Wang S, Zhang L, Mao Z, Tang P. A systematic review and meta-analysis of Ilizarov methods in the treatment of infected nonunion of tibia and femur. *PLoS One.* 2015;10(11):e0141973. Epub 2015 Nov 3.
48. Henderson DJ, Barron E, Hadland Y, Sharma HK. Functional outcomes after tibial shaft fractures treated using the Taylor Spatial Frame. *J Orthop Trauma.* 2015 Feb;29(2):e54-9.
49. Fletcher MD, Solomin LN. Definitive management of significant soft tissue loss associated with open diaphyseal fractures utilising circular external fixation without free tissue transfer, a comprehensive review of the literature and illustrative case. *Eur J Orthop Surg Traumatol.* 2015 Jan;25(1):65-75.

WHAT'S NEW IN LIMB LENGTHENING AND DEFORMITY CORRECTION

- 50.** Bernstein M, Fragomen AT, Sabharwal S, Barclay J, Rozbruch SR. Does integrated fixation provide benefit in the reconstruction of posttraumatic tibial bone defects? *Clin Orthop Relat Res.* 2015 Oct;473(10):3143-53.
- 51.** Varsegova TN, Shchudlo NA, Shchudlo MM, Saifutdinov MS, Stepanov MA. The effects of tibial fracture and Ilizarov osteosynthesis on the structural reorganization of sciatic and tibial nerves during the bone consolidation phase and after fixator removal. *Strategies Trauma Limb Reconstr.* 2015 Aug;10(2):87-94. Epub 2015 Aug 8.
- 52.** Fadel M, Ahmed MA, Al-Dars AM, Maabed MA, Shawki H. Ilizarov external fixation versus plate osteosynthesis in the management of extra-articular fractures of the distal tibia. *Int Orthop.* 2015 Mar;39(3):513-9. Epub 2014 Dec 5.
- 53.** El-Mowafi H, El-Hawary A, Kandil Y. The management of tibial pilon fractures with the Ilizarov fixator: the role of ankle arthroscopy. *Foot (Edinb).* 2015 Dec;25(4):238-43. Epub 2015 Aug 20.
- 54.** Danoff JR, Saifi C, Goodspeed DC, Reid JS. Outcome of 28 open pilon fractures with injury severity-based fixation. *Eur J Orthop Surg Traumatol.* 2015 Apr;25(3):569-75. Epub 2014 Sep 26.
- 55.** Yin P, Zhang L, Li T, Zhang L, Wang G, Li J, Liu J, Zhou J, Zhang Q, Tang P. Infected nonunion of tibia and femur treated by bone transport. *J Orthop Surg Res.* 2015;10:49. Epub 2015 Apr 10.
- 56.** Pallaro J, Angelliaume A, Dunet B, Lavoigne N, Tournier C, Fabre T. Reconstruction of femoral bone loss with a monoplane external fixator and bone transport. *Orthop Traumatol Surg Res.* 2015 Sep;101(5):583-7. Epub 2015 Jun 1.
- 57.** Azzam W, Atef A. Our experience in the management of segmental bone defects caused by gunshots. *Int Orthop.* 2016 Feb;40(2):233-8. Epub 2015 Jul 8.
- 58.** Grubor P, Milicevic S, Grubor M, Meccariello L. Treatment of bone defects in war wounds: retrospective study. *Med Arch.* 2015 Aug;69(4):260-4. Epub 2015 Aug 4.
- 59.** Tilkeridis K, Chari B, Cheema N, Tryfonidis M, Khaleel A. The Ilizarov method for the treatment of complex tibial fractures and non-unions in a mass casualty setting: the 2005 earthquake in Pakistan. *Strategies Trauma Limb Reconstr.* 2015 Apr;10(1):13-20. Epub 2015 Mar 13.
- 60.** Keightley AJ, Nawaz SZ, Jacob JT, Unnithan A, Elliott DS, Khaleel A. Ilizarov management of Schatzker IV to VI fractures of the tibial plateau: 105 fractures at a mean follow-up of 7.8 years. *Bone Joint J.* 2015 Dec;97-B(12):1693-7.
- 61.** Vollans S, Chaturvedi A, Sivasankaran K, Madhu T, Hadland Y, Allgar V, Sharma HK. Symptomatic venous thromboembolism following circular frame treatment for tibial fractures. *Injury.* 2015;46(6):1108-11. Epub 2015 Apr 15.
- 62.** Zhong W, Lu S, Chai Y, Wen G, Wang C, Han P. One-stage reconstruction of complex lower extremity deformity combining Ilizarov external fixation and sural neurocutaneous flap. *Ann Plast Surg.* 2015 Apr;74(4):479-83.
- 63.** Rozbruch SR, Rozbruch ES, Zonshayn S, Borst EW, Fragomen AT. What is the utility of a limb lengthening and reconstruction service in an academic department of orthopaedic surgery? *Clin Orthop Relat Res.* 2015 Oct;473(10):3124-32.
- 64.** W-Dahl A, Toksvig-Larsen S, Lindstrand A. Ten-year results of physical activity after high tibial osteotomy in patients with knee osteoarthritis. *Knee Surg Sports Traumatol Arthrosc.* 2015 Jul 14. [Epub ahead of print].
- 65.** Faschingbauer M, Nelitz M, Urlaub S, Reichel H, Dornacher D. Return to work and sporting activities after high tibial osteotomy. *Int Orthop.* 2015 Aug;39(8):1527-34. Epub 2015 Feb 26.
- 66.** Pitta M, Davis W 3rd, Argintar EH. Arthroscopic management of osteoarthritis. *J Am Acad Orthop Surg.* 2016 Feb;24(2):74-82.
- 67.** Bode G, Ogon P, Pestka J, Zwingmann J, Feucht M, Südkamp N, Niemeyer P. Clinical outcome and return to work following single-stage combined autologous chondrocyte implantation and high tibial osteotomy. *Int Orthop.* 2015 Apr;39(4):689-96. Epub 2014 Oct 10.
- 68.** Schuster P, Schulz M, Mayer P, Schlumberger M, Immendoerfer M, Richter J. Open-wedge high tibial osteotomy and combined abrasion/microfracture in severe medial osteoarthritis and varus malalignment: 5-year results and arthroscopic findings after 2 years. *Arthroscopy.* 2015 Jul;31(7):1279-88. Epub 2015 Apr 7.
- 69.** Schuster P, Schulz M, Richter J. Combined biplanar high tibial osteotomy, anterior cruciate ligament reconstruction, and abrasion/microfracture in severe medial osteoarthritis of unstable varus knees. *Arthroscopy.* 2016 Feb;32(2):283-92. Epub 2015 Sep 15.
- 70.** Stupina TA, Stepanov MA, Teplen'kii MP. Role of subchondral bone in the restoration of articular cartilage. *Bull Exp Biol Med.* 2015 Apr;158(6):820-3. Epub 2015 Apr 21.
- 71.** Lash NJ, Feller JA, Batty LM, Wasiak J, Richmond AK. Bone grafts and bone substitutes for opening-wedge osteotomies of the knee: a systematic review. *Arthroscopy.* 2015 Apr;31(4):720-30. Epub 2015 Jan 14.
- 72.** Cameron JJ, McCauley JC, Kermanshahi AY, Bugbee WD. Lateral opening-wedge distal femoral osteotomy: pain relief, functional improvement, and survivorship at 5 years. *Clin Orthop Relat Res.* 2015 Jun;473(6):2009-15. Epub 2014 Dec 24.
- 73.** Harris JD, Hussey K, Wilson H, Pilz K, Gupta AK, Gomoll A, Cole BJ. Biological knee reconstruction for combined malalignment, meniscal deficiency, and articular cartilage disease. *Arthroscopy.* 2015 Feb;31(2):275-82. Epub 2014 Oct 16.
- 74.** Konopka JF, Gomoll AH, Thornhill TS, Katz JN, Losina E. The cost-effectiveness of surgical treatment of medial unicompartmental knee osteoarthritis in younger patients: a computer model-based evaluation. *J Bone Joint Surg Am.* 2015 May 20;97(10):807-17.
- 75.** Burghardt RD, Manzotti A, Bhav A, Paley D, Herzenberg JE. Tibial lengthening over intramedullary nails: a matched case comparison with Ilizarov tibial lengthening. *Bone Joint Res.* 2016 Jan;5(1):1-10.
- 76.** Ryu KJ, Kim BH, Hwang JH, Kim HW, Lee DH. Reamed intramedullary nailing has an adverse effect on bone regeneration during the distraction phase in tibial lengthening. *Clin Orthop Relat Res.* 2016 Mar;474(3):816-24. Epub 2015 Oct 27.
- 77.** Dehghan MM, Baghaban Eslaminejad M, Motallebizadeh N, Ashrafi Halan J, Tagiyar L, Soroori S, Nikmahzar A, Pedram M, Shahverdi A, Kazemi Mehrjeri H, Izadi S. Transplantation of autologous bone marrow mesenchymal stem cells with platelet-rich plasma accelerate distraction osteogenesis in a canine model. *Cell J.* 2015 Summer;17(2):243-52. Epub 2015 Jul 11.
- 78.** Paley D. PRECICE intramedullary limb lengthening system. *Expert Rev Med Devices.* 2015 May;12(3):231-49. Epub 2015 Feb 18.
- 79.** Landge V, Shabtai L, Gesheff M, Specht SC, Herzenberg JE. Patient satisfaction after limb lengthening with internal and external devices. *J Surg Orthop Adv.* 2015 Fall;24(3):174-9.
- 80.** Horn J, Grimsrud Ø, Dagsgard AH, Huhnstock S, Steen H. Femoral lengthening with a motorized intramedullary nail. *Acta Orthop.* 2015 Apr;86(2):248-56. Epub 2014 Sep 5.
- 81.** Hawi N, Kenaway M, Panzica M, Stuebig T, Omar M, Krettek C, Lioudakis E. Nail-medullary canal ratio affects mechanical axis deviation during femoral lengthening with an intramedullary distractor. *Injury.* 2015 Nov;46(11):2258-62. Epub 2015 May 27.
- 82.** Bhav A, Shabtai L, Ong PH, Standard SC, Paley D, Herzenberg JE. Custom knee device for knee contractures after internal femoral lengthening. *Orthopedics.* 2015 Jul 1;38(7):e567-72.
- 83.** Elbatawy Y, Ragab IM. Safe cosmetic leg lengthening for short stature: long-term outcomes. *Orthopedics.* 2015 Jul 1;38(7):e552-60.
- 84.** Zampogna B, Vasta S, Amendola A, Uribe-Echevarria Marbach B, Gao Y, Papalia R, Denaro V. Assessing lower limb alignment: comparison of standard knee xray vs long leg view. *Iowa Orthop J.* 2015;35:49-54.
- 85.** Lee SA, Choi SH, Chang MJ. How accurate is anatomic limb alignment in predicting mechanical limb alignment after total knee arthroplasty? *BMC Musculoskelet Disord.* 2015;16:323. Epub 2015 Oct 27.
- 86.** Cho Y, Ko Y, Lee W. Relationships among foot position, lower limb alignment, and knee adduction moment in patients with degenerative knee osteoarthritis. *J Phys Ther Sci.* 2015 Jan;27(1):265-8. Epub 2015 Jan 9.
- 87.** Weinberg DS, Park PJ, Morris WZ, Liu RW. Femoral version and tibial torsion are not associated with hip or knee arthritis in a large osteological collection. *J Pediatr Orthop.* 2015 Jul 24. [Epub ahead of print].
- 88.** Weinberg DS, Liu RW. The association of tibia femur ratio and degenerative disease of the spine, hips, and knees. *J Pediatr Orthop.* 2015 Sep 20. [Epub ahead of print].
- 89.** Feldman DS, Goldstein RY, Kurland AM, Sheikh Taha AM. Intra-articular osteotomy for genu valgum in the knee with a lateral compartment deficiency. *J Bone Joint Surg Am.* 2016 Jan 20;98(2):100-7.
- 90.** Alexis F, Herzenberg JE, Nelson SC. Deformity correction in Haiti with the Taylor Spatial Frame. *Orthop Clin North Am.* 2015 Jan;46(1):9-19. Epub 2014 Oct 18.
- 91.** Morasiewicz P, Dragan S, Dragan SŁ, Wrzosek Z, Pawik Ł. Pedobarographic analysis of body weight distribution on the lower limbs and balance after Ilizarov corticotomies. *Clin Biomech (Bristol, Avon).* 2016 Jan;31:2-6. Epub 2015 Oct 23.
- 92.** Morasiewicz P, Konieczny G, Pawik Ł, Dragan S. Sport and physical activity in patients after derotational corticotomies with the Ilizarov method. *Acta Orthop Belg.* 2015 Mar;81(1):90-9.
- 93.** Kovoor CC, George VV, Jayakumar R, Guild AJ, Bhaskar D, Cryiac A. Total and subtotal amputation of lower limbs treated by acute shortening, revascularization and early limb lengthening with Ilizarov ring fixation - a retrospective study. *Injury.* 2015 Oct;46(10):1964-8. Epub 2015 Jul 23.
- 94.** Lam A, Garrison G, Rozbruch SR. Lengthening of tibia after trans-tibial amputation: use of a weight bearing external fixator-prosthesis composite. *HSS J.* 2016 Feb;12(1):85-90. Epub 2015 Sep 8.
- 95.** Felcher SM, Stinner DJ, Krueger CA, Wilken JM, Gajewski DA, Hsu JR; Skeletal Trauma Research Consortium (STRc). Falls in a young active amputee population: a frequent cause of rehospitalization? *Mil Med.* 2015 Oct;180(10):1083-6.
- 96.** Jauregui JJ, Bor N, Thakral R, Standard SC, Paley D, Herzenberg JE. Life- and limb-threatening infections following the use of an external fixator. *Bone Joint J.* 2015 Sep;97-B(9):1296-300.

WHAT'S NEW IN LIMB LENGTHENING AND DEFORMITY CORRECTION

- 97.** Ortiz D 3rd, Blair JA, Dromsky DM, Pyo J, Owens JG, Hsu JR; Skeletal Trauma Research Consortium (STReC). Collaborative establishment of an integrated orthotic and rehabilitation pathway. *J Surg Orthop Adv.* 2015 Fall;24(3):155-8.
- 98.** Rivera JC, Hsu JR, Noel SP, Wenke JC, Rathbone CR. Locally delivered nonsteroidal antiinflammatory drug: a potential option for heterotopic ossification prevention. *Clin Transl Sci.* 2015 Oct;8(5):591-3. Epub 2015 Jun 15.
- 99.** Barbier D, Neretin A, Journeau P, Popkov D. Gradual metatarsal lengthening by external fixation: a new classification of complications and a stable technique to minimize severe complications. *Foot Ankle Int.* 2015 Nov;36(11):1369-77. Epub 2015 Jul 10.
- 100.** Kawoosa AA, Baba MA, Wani IH, Khursheed O, Dar FA, Bashir A. Ankle arthrodesis using the Ilizarov technique in difficult situations - a prospective study with mid-to long-term follow up. *Ortop Traumatol Rehabil.* 2015 Mar-Apr;17(2):147-53.
- 101.** Rabinovich RV, Haleem AM, Rozbruch SR. Complex ankle arthrodesis: review of the literature. *World J Orthop.* 2015 Sep 18;6(8):602-13.
- 102.** Cavadas PC, Thione A. Reconstruction of the lateral malleolus in a type-Ib fibular hemimelia with a microvascular proximal fibular flap: a case report. *J Pediatr Orthop B.* 2015 Jul;24(4):370-2.
- 103.** Lohia LK, Meena S, Kanojia RK. Comparative study of complete subtalar release and Joshi's external stabilization system in the management of neglected and resistant idiopathic clubfoot. *Foot Ankle Surg.* 2015 Mar;21(1):16-21. Epub 2014 Aug 27.
- 104.** Sinkin JC, Reilly M, Cralley A, Kim PJ, Steinberg JS, Cooper P, Evans KK, Attinger CE. Multidisciplinary approach to soft-tissue reconstruction of the diabetic Charcot foot. *Plast Reconstr Surg.* 2015 Feb;135(2):611-6.
- 105.** Sala F, Catagni M, Pili D, Capitani P. Elbow arthrodesis for post-traumatic sequelae: surgical tactics using the Ilizarov frame. *J Shoulder Elbow Surg.* 2015 Nov;24(11):1757-63.
- 106.** Hamada Y, Sairoy K, Hibino N, Kobayashi A. Correction of severe contracture of intrinsic plus hand with a modified Ilizarov mini-fixator: correction with an Ilizarov mini-fixator for severe hand contracture. *J Hand Microsurg.* 2015 Jun;7(1):161-5. Epub 2014 May 14.
- 107.** Hamada Y, Kobayashi A, Sairoy K, Sato R, Hibino N. Correction of a hyperextension deformity at the metacarpophalangeal joint by arthroplasty for osteoarthritis of the thumb carpometacarpal joint followed by external fixator: a case series: modified Ilizarov method for correction of a collapsed thumb deformity due to carpometacarpal osteoarthritis. *J Hand Microsurg.* 2015 Jun;7(1):67-72. Epub 2015 Feb 28.
- 108.** Hamada Y, Hibino N, Kobayashi A. Surgical rehabilitation for correction of severe flexion contracture of the proximal interphalangeal joint by modified Ilizarov method. *J Hand Surg Eur Vol.* 2015 Feb;40(2):208-10. Epub 2013 Dec 11.
- 109.** Oh CS, Jung ST, Cho YJ, Ahn YS, Na BR. Bone transport for reconstruction in benign bone tumors. *Clin Orthop Surg.* 2015 Jun;7(2):248-53. Epub 2015 May 18.
- 110.** Borzunov DY, Balaev PI, Subramanyam KN. Reconstruction by bone transport after resection of benign tumors of tibia: a retrospective study of 38 patients. *Indian J Orthop.* 2015 Sep-Oct;49(5):516-22.
- 111.** Alzahrani MM, Rauch F, Hamdy RC. Does sclerostin depletion stimulate fracture healing in a mouse model? *Clin Orthop Relat Res.* 2016 May;474(5):1294-302. Epub 2015 Nov 25.
- 112.** Morcos MW, Al-Jallad H, Hamdy R. Comprehensive review of adipose stem cells and their implication in distraction osteogenesis and bone regeneration. *Biomed Res Int.* 2015;2015:842975. Epub 2015 Sep 13.
- 113.** Nayef L, Mekhail M, Benameur L, Rendon JS, Hamdy R, Tabrizian M. A combinatorial approach towards achieving an injectable, self-contained, phosphate-releasing scaffold for promoting biomineralization in critical size bone defects. *Acta Biomater.* 2016 Jan;29:389-97. Epub 2015 Oct 17.
- 114.** Grünwald L, Döbele S, Höntzsch D, Slongo T, Stöckle U, Freude T, Schröter S. Callus massage after distraction osteogenesis using the concept of lengthening then dynamic plating. *Strategies Trauma Limb Reconstr.* 2015 Aug;10(2):129-35. Epub 2015 Sep 4.
- 115.** Lundblad H, Maguire GQ Jr, Karlsson-Thur C, Jonsson C, Noz ME, Zeleznik MP, Jacobsson H, Weidenhielm L. Using PET/CT bone scan dynamic data to evaluate tibia remodeling when a Taylor Spatial Frame is used: short and longer term differences. *Biomed Res Int.* 2015;2015:574705. Epub 2015 Sep 7.
- 116.** Mora-Macías J, Reina-Romo E, López-Pliego M, Giráldez-Sánchez MA, Domínguez J. In vivo mechanical characterization of the distraction callus during bone consolidation. *Ann Biomed Eng.* 2015 Nov;43(11):2663-74. Epub 2015 May 9.
- 117.** Stogov MV, Tushina NV, Emanov AA. Serum concentration of growth factors in dogs under different conditions of distraction osteogenesis. *Bull Exp Biol Med.* 2015 Dec;160(2):213-5. Epub 2015 Dec 8.
- 118.** Bairaktari C, Athanassiou G, Panagiotopoulos E, Deligianni D. Towards a solution of the wires' slippage problem of the Ilizarov external fixator. *Eur J Orthop Surg Traumatol.* 2015 Apr;25(3):435-42. Epub 2014 Aug 20.
- 119.** Capanni F, Hansen K, Fitzpatrick DC, Madey SM, Bottlang M. Elastically suspending the screw holes of a locked osteosynthesis plate can dampen impact loads. *J Appl Biomech.* 2015 Jun;31(3):164-9. Epub 2015 Feb 2.
- 120.** Henderson DJ, Rushbrook JL, Stewart TD, Harwood PJ. What are the biomechanical effects of half-pin and fine-wire configurations on fracture site movement in circular frames? *Clin Orthop Relat Res.* 2016 Apr;474(4):1041-9. Epub 2015 Dec 7.
- 121.** Jabbar Y, Khaleel A. Experimental model for controlling shear using the Ilizarov frame. *Clin Biomech (Bristol, Avon).* 2015 Nov;30(9):995-1001. Epub 2015 Jul 11.
- 122.** Skomoroshko PV, Vilensky VA, Hammouda AI, Fletcher MD, Solomin LN. Mechanical rigidity of the Ortho-SUV frame compared to the Ilizarov frame in the correction of femoral deformity. *Strategies Trauma Limb Reconstr.* 2015 Apr;10(1):5-11. Epub 2015 Feb 26.
- 123.** Kim KN, Hong JP, Park SW, Kim SW, Yoon CS. Overcoming the obstacles of the Ilizarov device in extremity reconstruction: usefulness of the perforator as the recipient vessel. *J Reconstr Microsurg.* 2015 Jul;31(6):420-5. Epub 2015 Mar 31.
- 124.** Lowery K, Dearden P, Sherman K, Mahadevan V, Sharma H. Cadaveric analysis of capsular attachments of the distal femur related to pin and wire placement. *Injury.* 2015;46(6):970-4. Epub 2015 Mar 16.
- 125.** Dearden P, Lowery K, Sherman K, Mahadevan V, Sharma H. Fibular head transfixion wire and its relationship to common peroneal nerve: cadaveric analysis. *Strategies Trauma Limb Reconstr.* 2015 Aug;10(2):73-8. Epub 2015 May 28.
- 126.** Ktistakis I, Guerado E, Giannoudis PV. Pin-site care: can we reduce the incidence of infections? *Injury.* 2015 Sep;46(Suppl 3):S35-9.